(19) 世界知的所有権機関 国際事務局



. I COLO DE REPORTE EL BORRO DE LO COLO COLO DE LA COLO

(43) 国際公開日 2003 年1 月30 日 (30.01.2003)

PCT

(10) 国際公開番号 WO 03/007819 A1

(51) 国際特許分類7:

A61B 5/15

(21) 国際出願番号:

PCT/JP02/07333

(22) 国際出願日:

2002年7月18日 (18.07.2002)

(25) 国際出願の言語:

日本語

(26) 国際公開の言語:

日本語

(30) 優先権データ:

特願2001-220259 2001年7月19日(19.07.2001) JP 特願2001-236976 2001年8月3日(03.08.2001) JP

- (71) 出願人 (米国を除く全ての指定国について): アークレイ株式会社 (ARKRAY, INC.) [JP/JP]; 〒601-8045 京都府 京都市 南区東九条西明田町 5 7 Kyoto (JP)
- (72) 発明者; および
- (75) 発明者/出願人 (米国についてのみ): 坂田 哲也 (SAKATA,Tetsuya) [JP/JP]; 〒601-8045 京都府 京都 市 南区東九条西明田町 5 7 アークレイ株式会社 内 Kyoto (JP). 松本 大輔 (MATSUMOTO,Daisuke)


[JP/JP]; 〒601-8045 京都府 京都市 南区東九条西明田町57 アークレイ株式会社内 Kyoto (JP). 笠井 督夫(KASAI,Tokuo) [JP/JP]; 〒601-8045 京都府 京都市 南区東九条西明田町57 アークレイ株式会社内 Kyoto (JP)

- (74) 代理人: 吉田 稔, 外(YOSHIDA, Minoru et al.); 〒 543-0014 大阪府 大阪市 天王寺区玉造元町 2 番 3 2-1 3 0 1 Osaka (JP).
- (81) 指定国 (国内): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) 指定国 (広域): ARIPO 特許 (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), ユーラシア特許 (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), ヨーロッパ 特許 (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,

/続葉有/

(54) Title: PIERCING DEVICE

(54) 発明の名称: 穿刺装置



(57) Abstract: A piercing device (A), comprising a piercing body (3) allowed to move forward in a first direction from the inner bottom part of a housing (2) to the tip end of a tubular body (8) and an analysis part (4) disposed in the tubular body (8), wherein the analysis part (4) is formed movable in a second direction when receiving a force in a second direction opposite to the first direction, and when skin (S) is raised, the analysis part (4) is moved in the second direction by following the rise of the skin to properly lead blood (b) from the skin (S) to the analysis part (4).

FIG.1




FIG.2




FIG.3




FIG.4




FIG.5




FIG.6




FIG.7







FIG.8






7/19

FIG.10



WO 03/007819



WO 03/007819




FIG.13




FIG.14




FIG.15




FIG.16




FIG.17




FIG.18




FIG.19




FIG.20




FIG.21




FIG.22

